



Botox Pre-Treatment Information

Botox, a neurotoxin produced by the bacterium *Clostridium A*. Botox, can relax the muscles on areas of the face and neck that cause wrinkles associated with facial expressions. Treatment with Botox can cause your facial expression line or wrinkles to essentially disappear. Areas most frequently treated are: a) glabellar area of frown lines, located between the eyes; b) crows feet (lateral areas of the eyes); and c) forehead wrinkles. Botox is injected into the muscles with a very thin needle and is almost painless. You may feel a slight burning sensation while the solution is being injected. The procedure takes about 15-20 minutes and the results last 3-4 months. With repeated treatments, the results may last longer.

These are some contraindication to having treatment:

- Allergies to eggs, egg products, albumin.
- Significant autoimmune or neurological diseases.
- Currently taking aminoglycoside antibiotics.
- Pregnant or lactating.

These are some possible side effects:

- Some initial swelling, pain, itching, discoloration, bruising or tenderness at the injection site. You could experience increased bruising or bleeding at the injection site if you are on aspirin or other non-steroidal anti-inflammatory drugs such as Advil. These reactions generally lessen or disappear within a few days but may last for a week or longer.
- As with all injections, this procedure carries the risk of infection.

Additional Information:

- Individual results may vary.
- You can purchase Arnica over-the-counter at the drug store to help alleviate bruising.
- For medical reasons, photographs will be taken and documented in your confidential patient file.

Please call if you have any questions prior to your appointment.