



Halo Hybrid Fractional Laser Pre Treatment Considerations

- Arrive 45 minutes prior to your scheduled procedure for topical numbing and pre-treatment photos.
- If you have previously suffered from facial cold sores, there is a risk that this treatment could contribute to a recurrence. It is recommended that you begin antiviral therapy one day prior to your procedure.
- No one who has taken the medication Accutane or its generic forms within the last year may have this procedure.
- No one on anti-coagulants may have this procedure.
- Hydroquinone (Obagi Skin Bleaching and Corrector Cream, HQ 4%) should be started 4 weeks prior to procedure for Fitzpatrick Skin Types 3+, ethnic skin, melasma patients, or anyone with a history of hyperpigmentation or pigmentation concerns.
* For those with a history or concern for extreme hyperpigmentation, 7% HQ should be used.
- Retinol products should be discontinued one week prior to Halo procedure
- No direct or intense sun exposure should occur one week prior to or following laser treatment
- No spray tan/self-tanning products should be used starting 2 weeks prior to the procedure
- Recommended Products for Optimal Halo Results:
 - ***SkinMedica Facial Cleanser***
 - ***TNS Recovery Complex***
 - ***Essential Defense SPF35 Post Treatment Sunscreen***
 - ***Ultrasheer Moisturizer or TNS Ceramide Treatment Cream***

*Lytera 2.0 highly recommended for long term skin lightener, post treatment.

Please call should you have any questions prior to your appointment.

Tel:

Tel: 619.293.7233
revive.md